

The leaf essential oils of *Picea chihuahuana* Martinez and *P. martinezii* Patterson (Pinaceae)**Robert P. Adams**

Biology Department, Baylor University, Box 97388, Waco, TX 76798, USA Robert_Adams@baylor.edu

and

Frank Callahan

P. O. Box 5531, Central Point, OR 97502

ABSTRACT

The first comprehensive analysis of the volatile leaf terpenoids of *Picea chihuahuana* and *P. martinezii* is reported. The volatile leaf oil of *P. chihuahuana* is dominated by bornyl acetate (47.9%) with moderate amounts of α -pinene (4.7), camphene (6.3), β -pinene (5.2), camphene (6.2), limonene (7.4) and β -phellandrene (7.3%). The leaf oil of *P. martinezii* is dominated by bornyl acetate (26.4%), α -pinene (16.6) and myrcene (15.1%), with moderate amounts of camphene (5.6) and β -pinene (9.7%). For such closely related species (Patterson and Harrod, 1994), the oils are quite differentiated. Components unique in *P. chihuahuana* oil are: α -phellandrene, cis- and trans-p-menth-2-en-1-ol, cis- and trans-piperitol, piperitone, cis- and trans-piperitol acetate, methyl isopimarate, methyl levopimarate and methyl abietate. Unique compounds in *P. martinezii* oil include: α -muurolene, δ -cadinene, caryophyllene oxide, humulene epoxide II, epi- α -cadinol, epi- α -muurolol, α -muurolol, α -cadinol, germacrene-4(15),5,10(14)-triene-1-al and neo-abienal. Published on-line www.phytologia.org *Phytologia* 96(4): 260-263 (Oct 1, 2014). ISSN 030319430

KEY WORDS: *Picea chihuahuana*, *P. martinezii*, terpene composition, volatile leaf oils, Pinaceae.

Taylor, Patterson and Harrod (1994) used a combination of morphology, phenolics and terpenoids to examine Mexican species of *Picea*. Of interest to the present study, was the observation that *Picea chihuahuana* and *P. martinezii* were shown to be distinct species by their morphology (Fig. 2), phenolics (Fig. 5) and terpenoids (Fig. 6), re-confirming their previous study (Taylor and Patterson, 1980). Taylor, Patterson and Harrod (1994) listed α -pinene, camphene, β -pinene, myrcene, γ -terpinene, piperitone, citronellol, geraniol, camphor, and bornyl acetate as present in six species of *Picea* examined, but did not give any composition data.

Recently, Jaramillo-Correa et al. (2006) examined mt and cp DNA markers among *P. chihuahuana* and *P. martinezii* and concluded that they are not conspecific, but distinct species.

Aside from the qualitative report of Taylor, Patterson and Harrod (1994), no reports have been published on the composition of the leaf volatile oils of *Picea chihuahuana* or *P. martinezii*. The purpose of the present paper is to give the first detailed report on the composition of the volatile oils of these species.

MATERIALS AND METHODS

Leaf samples were collected from *Picea chihuahuana* (Adams 14329-14331) from trees cultivated at Medford city park, Medford, OR (grown from seed collected on Rio Oteros, near Creel, Chihuahua, Mexico) and from *P. martinezii* (Adams 14326-14328) cultivated at Frank Callahan's land, Central Point, OR, grown from seed collected by Frank Callahan in Nuevo Leon, Mexico. Voucher specimens are deposited in the herbarium, Baylor University.

Fresh, frozen leaves (200 g) were steam distilled for 2 h using a circulatory Clevenger-type apparatus (Adams, 1991). The oil samples were concentrated (ether trap removed) with nitrogen and the samples stored at -20°C until analyzed. The extracted leaves were oven dried (100°C, 48 h) for determination of oil yields.

The oils were analyzed on a HP5971 MSD mass spectrometer, scan time 1/ sec., directly coupled to a HP 5890 gas chromatograph, using a J & W DB-5, 0.26 mm x 30 m, 0.25 micron coating thickness, fused silica capillary column (see Adams, 2007 for operating details). Identifications were made by library searches of our volatile oil library (Adams, 2007), using the HP Chemstation library search routines, coupled with retention time data of authentic reference compounds. Quantitation was by FID on an HP 5890 gas chromatograph using a J & W DB-5, 0.26 mm x 30 m, 0.25 micron coating thickness, fused silica capillary column using the HP Chemstation software.

RESULTS AND DISCUSSION

The leaf oils of *P. chihuahuana* and *P. martinezii* were clear in color and the yields (w/DW) very small: 0.34% and 0.25%, respectively. The volatile leaf oil of *P. chihuahuana* is dominated by bornyl acetate (47.9%, Table 1) with moderate amounts of α -pinene (4.7), camphene (6.3), β -pinene (5.2), camphene (6.2), limonene (7.4) and β -phellandrene (7.3%). The leaf oil of *P. martinezii* is dominated by bornyl acetate (26.4%, Table 1), α -pinene (16.6) and myrcene (15.1%) with moderate amounts of camphene (5.6) and β -pinene (9.7%). For such closely related species (Patterson and Harrod, 1994), the oils are quite different (Table 1). There are several unique oil components in *P. chihuahuana*: α -phellandrene, cis- and trans-p-menth-2-en-1-ol, cis- and trans-piperitol, piperitone, cis- and trans-piperitol acetate, methyl isopimarate, methyl levopimarate and methyl abietate. Unique oil compounds in *P. martinezii* include: α -muurolene, δ -cadinene, caryophyllene oxide, humulene epoxide II, epi- α -cadinol, epi- α -muurolol, α -muurolol, α -cadinol, germacra-4(15),5,10(14)-triene-1-al and neo-abienal (Table 1).

The leaf volatile terpenoids data support the conclusions of Patterson and Harrod (1994) and Jaramillo-Correa et al. (2006) that *P. chihuahuana* and *P. martinezii* are distinct species.

It might be noted that *P. chihuahuana*, in cultivation in Medford, OR (Fig. 1), is a robust, fast growing tree that will likely be widely cultivated for its glaucous foliage, shape and hardness.

ACKNOWLEDGEMENTS

This research supported with funds from Baylor University.



Figure 1. *Picea chihuahuana*, seed cone (insert) in cultivation at the Medford City Park, OR with Frank Callahan.

LITERATURE CITED

- Adams, R. P. 1991. Cedarwood oil - Analysis and properties. pp. 159-173. in: Modern Methods of Plant Analysis, New Series: Oil and Waxes. H.-F. Linskens and J. F. Jackson, eds. Springer- Verlag, Berlin.
- Adams, R. P. 2007. Identification of essential oil components by gas chromatography/ mass spectrometry. 4th ed. Allured Publ., Carol Stream, IL.
- Jaramillo-Correa, J. Beaulieu, F. T. Ledig and J. Bousquet. Decoupled mitochondrial and chloroplast DNA population structure reveals Holocene collapse and population isolation in a threatened Mexican-endemic conifer. Molec. Ecol. 15: 2787-2800.
- Taylor, R. J., T. F. Patterson and R. J. Harrod. 1994. Systematics of Mexico spruces - revisited. Syst. Bot. 19: 47-89.
- Taylor, R. J. and T. F. Patterson. 1980. Biosystematics of Mexican spruce species. Taxon 29: 421-469.

Table 1. Comparison of leaf oil compositions of *Picea chihuahuana* and *P. martinezii*. Compounds in bold face appear to separate the taxa. Compositional values less than 0.1% are denoted as traces (t). Unidentified components less than 0.5% are not reported. KI is the Kovat's Index using a linear calculation on DB-5 column.

| KI | compound | <i>chihuahuana</i> | <i>martinezii</i> |
|-------------|---|--------------------|-------------------|
| 846 | (E)-hexenal | 0.2 | 0.3 |
| 921 | tricyclene | 0.4 | 0.4 |
| 932 | α-pinene | 4.7 | 16.6 |
| 946 | camphene | 6.3 | 5.6 |
| 969 | sabinene | t | t |
| 974 | β-pinene | 5.2 | 9.7 |
| 988 | myrcene | 3.2 | 15.1 |
| 997 | ethyl hexanoate | 0.2 | - |
| 1002 | α-phellandrene | 0.7 | - |
| 1008 | δ -3-carene | t | - |
| 1014 | α -terpinene | 0.1 | - |
| 1020 | p-cymene | t | t |
| 1024 | limonene | 7.4 | 1.6 |
| 1025 | β-phellandrene | 7.3 | 2.4 |
| 1032 | (Z)- β -ocimene | t | - |
| 1054 | γ -terpinene | t | t |
| 1086 | terpinolene | 1.2 | 1.2 |
| 1095 | linalool | 0.3 | 0.2 |
| 1118 | cis-p-menth-2-en-1-ol | 0.9 | - |
| 1122 | α -campholenal | 0.1 | 0.3 |
| 1136 | trans-p-menth-2-en-1-ol | 0.7 | - |
| 1136 | trans-pinocarveol | - | 0.1 |
| 1141 | camphor | t | 0.1 |
| 1145 | camphene hydrate | 0.5 | 0.2 |
| 1165 | borneol | 1.7 | 0.5 |
| 1174 | terpinen-4-ol | 0.1 | 0.2 |
| 1186 | α -terpineol | 1.1 | 1.1 |
| 1190 | methyl salicylate | - | t |
| 1195 | cis-piperitol | 0.3 | - |
| KI | compound | <i>chihuahuana</i> | <i>martinezii</i> |
| 1207 | trans-piperitol | 0.6 | - |

| KI | compound | <i>chihuahuana</i> | <i>martinezii</i> |
|-------------|--|--------------------|-------------------|
| 1218 | endo-fenchyl acetate | t | - |
| 1223 | citronellol | t | - |
| 1249 | piperitone | 0.6 | - |
| 1287 | bornyl acetate | 47.9 | 26.4 |
| 1298 | trans-pinocarvyl acetate | 0.4 | 0.3 |
| 1324 | myrtenyl acetate | 0.1 | - |
| 1332 | cis-piperitol acetate | 0.1 | - |
| 1343 | trans-piperitol acetate | 0.2 | - |
| 1350 | α -longipinene | 0.1 | - |
| 1379 | geranyl acetate | - | 0.1 |
| 1396 | duvalene acetate | t | 0.1 |
| 1417 | (E)-caryophyllene | 0.3 | 2.1 |
| 1454 | (E)- β -farnesene | t | - |
| 1469 | n-dodecanol | t | - |
| 1480 | germacrene D | t | 1.1 |
| 1500 | α-muurolene | - | 0.5 |
| 1522 | δ-cadinene | - | 0.2 |
| 1565 | dodecanoic acid | t | - |
| 1582 | caryophyllene oxide | - | 0.3 |
| 1608 | humulene epoxide II | - | 0.3 |
| 1638 | epi-α-cadinol | - | 0.2 |
| 1640 | epi-α-muurolol | - | 0.2 |
| 1644 | α-muurolol | - | 0.1 |
| 1652 | α-cadinol | - | 0.8 |
| 1685 | germacra-4(15),5,10(14)-triene-1-al | - | 0.3 |
| 1959 | hexadecanoic acid | 0.2 | 0.5 |
| 1987 | iso-pimara-7,15-diene | 0.3 | 0.2 |
| 2182 | hexadecanoic acid, butyl ester | t | - |
| 2222 | abietal isomer | 0.3 | 0.9 |
| 2298 | methyl isopimarate | 0.2 | - |
| 2300 | tricosane | 0.2 | 0.4 |
| 2306 | methyl levopimarate | 0.4 | - |
| 2313 | abietal | 0.5 | 0.6 |
| 2375 | neo-abienal? | - | 0.4 |
| 2385 | methyl abietate | 0.1 | - |